

Application of 3D QSAR and Docking Studies in Optimization of Perylene diimides as Anti Cancer Agent

Catna Nagaraj Hemalatha¹, Vijey Aanandhi Muthukumar^{*2}

¹Department of Pharmaceutical Chemistry, School of Pharmaceutical Sciences, Vels Institute of Science, Technology and Advanced Studies, VISTAS, Pallavaram, Chennai, Tamilnadu, INDIA.

²Department of Pharmaceutical Chemistry and Analysis, School of Pharmaceutical Sciences, Vels Institute of Science, Technology and Advanced Studies, VISTAS, Pallavaram, Chennai, Tamilnadu, INDIA.

ABSTRACT

Introduction: Telomerase is an enzyme which binds to telomeres and increases its length which leads to extension of lifespan of cells. These enzymes are expressed at detectable levels in cancer cells which makes an attractive target for cancer therapy. The G Quadruplex ligands which bind to telomerase with respect to duplex genomic DNA is of special importance. The Perylene di imides are selected, designed and QSAR study has been done, finally from the QSAR results' docking has been done by G4LDB database. To compare and to narrow down the Docking results from G4LDB database, we have chosen AutoDock tool by selecting a target Telomerase protein (PDB ID: 4B18) to analyse the binding affinity of the protein with respect to the Perylene diimides. The best scored compounds will be efficient for designing new molecules as well as for the anti-cancer therapy. **Material:** G4LDB Database, AutoDock 4.2, Discovery Studio Visualizer 4.1. **Methods:** The study was to investigate and compare the results from G4LDB database with the AutoDock results for anti cancer activity of Perylene di imides. The results are visualized by Discovery Studio 4.1 Visualizer. Compound 20 and compound 48 shows best ligand interaction with the selected targets from G4LDB database. From the AutoDock results the compounds are docked with the specific Telomerase protein 4B18 and Compound 11 shows good binding energy when compared with the PIPER. **Results:** Compounds 11, 20 and 48 showed good biological activity also possessing best binding affinity with the target. **Discussion:** From the QSAR and Docking studies (G4LDB and AutoDock), 3 compounds (11, 20 and 48) showed good biological activity possessing a strong correlation coefficient, endorses that Perylene derivatives are having strong affinity with the targets. Docking has been done from the results of QSAR study, targeting Telomerase protein to study the binding affinity with the target. **Conclusions:** From the results, the best compounds will be efficient to inhibit telomerase enzyme and these compounds can be used to design new molecules which will be effective for anti-cancer therapy.

Key words: Perylene Derivatives, QSAR Plus, G-Quadruplex Ligand Database, Docking.

Submission Date: 09-08-2017;

Revision Date: 05-12-2017;

Accepted Date: 17-05-2018

DOI: 10.5530/ijper.52.4.77

Correspondence:

Dr. M. Vijey Aanandhi

Department of Pharmaceutical Chemistry and Analysis, School of Pharmaceutical Sciences, Vels Institute of Science, Technology and Advanced Studies, VISTAS, Pallavaram, Chennai, Tamil Nadu, INDIA.

Phone: +919840959519

E-mail: hodpchemistry@

velsuniv.ac.in

INTRODUCTION

Telomerase enzyme and its related proteins are having high expression in most type of tumour cells and it has been touted as most efficient and potent for cancer therapy. The compounds which interact or inhibit these proteins are Telomerase Inhibitors. The Guanine rich Quadruplex DNA (G4)

which is present in telomeric DNA stabilizes and affects the gene promoters, in turn disrupts the biological process. Perylene di imides are a promising class of G-Quadruplex Ligands which stabilize the G-Quadruplex structure, by stacking on the terminal G-Tetrads. In our previous



www.ijper.org

study we demonstrated the structural activity relationship of Observed and predicted Biological activity and it has been done in QSARPlus Module in VLifeMolecular Design Suite and performed docking study by using G4LDB Database (Guanine Rich Ligand Database). This QSAR module facilitates evaluation of descriptors and generates QSAR equation for predicting biological activity of new molecules.¹⁻² Based upon the study, new compounds are designed as the results shows the presence of steric descriptors in benzene ring with positive coefficient indicated the importance of steric interactions, bulky group can substitute and the presence of electrostatic descriptors with positive coefficient near to the bay region, while negative coefficient near to the imide indicated that electro negative groups should be substitutes on imide ring and the appearance of descriptors H-Bond acceptor with a positive coefficient suggest that an increased activity of the compound. These are significant in developing the novel perylene di imide derivatives.³ Further from the QSAR results, Docking has been done in G4LDB Database, is a unique and comprehensive database which compiles dataset of 28 G-Quadruplex Complex structures/targets are included in the database which was obtained from Protein Data Bank (PDB).⁴ In this study, to compare the results of Docking, we had done docking in AutoDock 4.2 to narrow down the results. From the QSAR and Docking results of G4LDB 9 compounds have been chosen for the study. The target selected for this study was a Telomerase protein (4B18:PDB ID) and by using this as target, docked the nine compounds to get to know the binding energy, ligand efficiency and the number of hydrogen bonds interacted with the target.⁵

MATERIALS AND METHODS

Molecular Docking was performed by the AutoDock 4.2 Tools. It helps to understand the binding properties in a steady state environment. In this study we used 4B18-a telomerase protein increases the binding affinity for importin proteins and promotes nuclear import of hTERT.

AutoDock calculations are performed in several steps: i) Preparation of coordinate files using AutoDock Tools, ii) Pre-calculation of atomic affinities using Auto Grid, iii) Docking of Ligands using AutoDock, and iv) Analysis of results using AutoDock Tools, Discovery Studio 4.1 Visualizer to visualize the interaction studies.⁶

preparation of ligands for docking

Perylene di imide ligands were taken. The compounds/ligands are the results of QSAR and G4LDB Database.

Accelrys Draw 4.2 was used to draw the structures and to convert 2D to 3D. All ligands are converted to PDB Format by using Discovery studio Visualizer which was used for coordinate files, which includes atomic partial charges and atom types. Torsion angles were calculated to assign the fixable and non-bonded rotations of the molecule.

Preparation of Protein (4B18)

High resolution crystallographic structures were retrieved from RCSB Protein Data Bank (www.rcsb.org). To understand the biological activity of a drug molecule as prototype therapeutic agent, the knowledge of binding selectivity towards protein environment is every essential. Docking study was performed to understand and correlate their biological efficacy towards the selected binding domain of the protein. We used the AutoDock 4.2 MGL tools version 1.5.6 software packages for the molecular docking experiment and Discovery Studio Visualizer to visualize the results. Polar Hydrogen atoms were added to the protein. The deletion of the both water molecules and the inorganic charges were done to avoid error. Gasteiger Charge of the macromolecule was added. Ligand docking was carried out applying the Lamarckian Genetic Algorithm (LGA) implemented in recognize the binding site. The grid size and spacing was set to recognize the binding site. The lowest binding energy conformers were selected out of different conformers. Other miscellaneous parameters were assigned to the default values obtained from AutoDock 4.2 program.

Molecular Docking Analysis was done by the output file of the docking that was generated after the study. The binding energy, Inhibition Constant, and the number of hydrogen bonds were considered for the analysis. Compound 11, 20 and 48 showed potent binding interactions with the protein 4B18 and compared with the standard PIPER. The visualization of the 3 best compounds are visualized by Discovery Studio Visualizer and the ligand interactions with the target, the distance between the ligand interactions are marked and the results are showed in **Figure a – p** for the compound – 11, 20, 48 and PIPER. The protein 4B18 was modelled by using PROCHECK, open source server and the Ramachandra plot for the protein was studied. The best compounds are compared with the standard PIPER and the binding energy values are showing more potent than the standard PIPER. The binding energy values for the compound 11, 20, 48 and PIPER are **-6.23, -6.42, -6.59 and -5.59 kcal/mol** respectively.

RESULTS AND CONCLUSION

QSAR study has been carried out by V Life MDS Software- QSAR Plus Module to predict and compare the biological activity of standard and newly designed compounds. QSAR has been done by developing variable regression methods. The compounds are divided into training and test set compounds by manual selection, random and sphere exclusion methods.⁷ The models are developed and based on regression values; we selected eight equations to design new compounds of perylene di imides. Out of 497 compounds, 59 compounds possess better biological activity when compared with the standard compounds.

These 59 selected compounds from QSAR study has been chosen for docking study. Docking has been done by G-Quadruplex Ligand Database (G4LDB).^{8,9} This is an online database which was having in built tools and performed by Open Babel 2.3.0 to predict the binding affinity with the targets. The targets (1LIH, 1NZM, 3CE5, 3SC8 and 2HRI) for the docking are selected based upon the literature survey and the selected compounds are docked. The results are visualized by Discovery studio Visualizer 4.1 Visualizer and the results of QSAR and G4LDB docked are tabulated in Table 1 and 2. From the results 9 compounds are selected and the results of these compounds are visualized using Discovery Studio 4.1 Visualizer. To narrow down the results, 2 best compounds are selected, and the compounds are Compound 20 and Compound 48. The hydrogen bond interactions and binding free energy levels, pK_i values are compared with the standard PIPER compound.^{10,11}





From the results of G4LDB Database, the 9 compounds are docked with the specific protein (PDB ID: 4B18), a telomerase protein. Docking has been done by Auto Dock 4.2. Finally, from the AutoDock results, Compound 11 shows good binding energy when compared with the standard PIPER compound. The study states as from the G4LDB Database 2 compounds possess good binding affinity, and from the Auto Dock **Compound 11** possess good binding energy. Based on the results from QSAR, G4LDB and AutoDock results compounds 11, 20 and 48 may have potent anti-cancer activity and from these compounds scaffolds we can design schemes for synthesis. In future studies we can use these scaffolds for experimental designing work for *in vitro* as well for *in vivo* methods.

The best potent compounds are studied to know the interactions with the nearby aminoacids and the ligand interactions are visualized by online PROCHECK server^{12,13} and the figures for the compounds are

Table 1: Newly designed compounds selected based upon the standard regression values- QSAR Plus Module- Vlife Sciences MDS Suite.

| CMP | QSAR No | Compounds | EQUA 1 | EQUA 2 | EQUA 3 | EQUA 4 | EQUA 5 | EQUA 6 | EQUA 7 | EQUA 8 |
|-----|---------|-------------------------------|---------|---------|---------|---------|---------|---------|---------|---------|
| 7 | 118 | cp_12_h_153_3D_opt.mol2 | 6.80698 | 5.30356 | 6.70725 | 5.32331 | 5.55316 | 7.23985 | 7.14242 | 7.22239 |
| 11 | 137 | cp_14a_a_172_3D_opt.mol2 | 6.44785 | 5.46056 | 5.20119 | 6.30898 | 6.50954 | 5.9222 | 5.76533 | 5.90495 |
| 20 | 213 | cp_daper_3c_b_234_3D_opt.mol2 | 6.87129 | 5.22616 | 5.25692 | 5.51287 | 6.52906 | 5.7838 | 5.35173 | 5.76984 |
| 21 | 214 | cp_daper_3c_c_249_3D_opt.mol2 | 7.01274 | 5.22621 | 5.25692 | 5.50151 | 6.47947 | 5.79134 | 5.53639 | 5.7771 |
| 30 | 326 | cp_piper_6_h_12_3D_opt.mol2 | 6.299 | 6.3151 | 5.2014 | 6.30537 | 6.40592 | 5.78281 | 5.66593 | 5.76909 |
| 33 | 339 | cp_pm2_cd_26_3D_opt.mol2 | 6.12263 | 5.63933 | 5.20144 | 5.10228 | 6.18314 | 5.80597 | 5.69873 | 5.79144 |
| 42 | 412 | cp_pol_6_d_98_3D_opt.mol2 | 6.27018 | 6.28015 | 6.70605 | 5.47861 | 6.04312 | 7.21727 | 7.08722 | 7.20086 |
| 48 | 426 | cp_pol_6_r_112_3D_opt.mol2 | 6.1255 | 6.23087 | 6.79506 | 5.47588 | 5.12905 | 7.22012 | 7.09411 | 7.20373 |
| 55 | 481 | cp_tel_c_167_3D_opt.mol2 | 8.29992 | 6.42848 | 5.25701 | 5.63671 | 6.62714 | 5.19589 | 5.85943 | 5.2165 |
| STD | Std | Compound PIPER 1 | 5.21084 | 6.24833 | 7.14924 | 6.31639 | 6.22921 | 6.43748 | 6.00598 | 6.40752 |

Table 2: pKi values – G-Quadruplex Ligand Database.

| S. No | Compounds | | 1L1H | 1NZM | 3CE5 | 3SC8 | 2HRI |
|-------|-----------|---|------|-------|------|------|------|
| 1 | CMP_11 |  | 7.33 | 12.4 | 8.34 | 6.5 | 4.26 |
| 2 | CMP_20 |  | 5.01 | 12.41 | 8.57 | 7.21 | 6.01 |
| 7 | CMP_48 |  | 8.66 | 12.1 | 8.54 | 6.65 | 6.6 |
| 9 | PIPER |  | 9.99 | 13.39 | 9.3 | 7.86 | 8.07 |




Figure 1: a) Binding Energy: -6.23 kcal/mol.




Figure 1: d) Hydrogen Bond Acceptor and Donor Representation of Protein with the Ligand.




Figure 1: b and c) Ligand Interactions with the nearby amino acids.




Figure 1: e) Binding Energy: -6.42 kcal/mol




Figure 1: f and g) Ligand Interactions with the nearby amino acids.




Figure 1: f and g) Ligand Interactions with the nearby amino acids.




Figure 1: j) Ligand Interactions with the nearby amino acids.




Figure 1: h) Hydrogen Bond Acceptor and Donor Representation of Protein with the Ligand.




Figure 1: k) Ligand Interactions with the nearby amino acids.




Figure 1: i) Binding Energy: -6.59 kcal/mol.




Figure 1: l) Hydrogen Bond Acceptor and Donor Representation of Protein with the Ligand.




Figure 1: m) Binding Energy: -5.59 kcal/mol.




Figure 1: n) Ligand Interactions with the nearby amino acids.




Figure 1: o) Ligand Interactions with the nearby amino acids.




Figure 1: p) Hydrogen Bond Acceptor and Donor Representation of Protein with the Ligand.




Figure 1: q) Ramachandra Plot for the protein 4B18.

shown respectively in Figure c(Compound 11), Figure g(compound 48), Figure k (Compound 48) and for Standard PIPER, Figure o. The Ramachandra plot for the protein was shown in Figure q.

DISCUSSION

Telomerase which is responsible for maintaining the length of the Telomere and scientific evidences proves it exhibits in tumor cells, and we chosen G-Quadruplex ligand-Perylene diimide, to study the biological activity and Binding affinity with respect to the targets. The standard compounds of Perylene diimides along with

their biological activity and descriptors are studied. From the QSAR results, we analysed as the perylene diimide derivatives are possessing the biological activity with the presence of steric descriptors in benzene ring with positive coefficients which indicates the importance of steric interactions, also bulky groups can substitute and the presence of electrostatic descriptors with positive coefficient near to the bay region, while negative coefficients near to the imide ring indicates that electro negative groups should be substituted on imide ring and the appearance of descriptors H-bond acceptor with a positive coefficient suggest that an increased activity of the compound. These are significant in developing the novel perylene diimide derivatives. Based upon these results, we have selected 59 compounds from 491 newly designed compounds, where the selection was based on the biological activity (r^2) ranging from 5 to 9 from all the 8 equations for the synthesis.

From the results of QSAR, Docking has been done by G4LDB Database and Autodock and the PDB ID for G4LDB database are 1LIH, 1NZM, 3CE5, 3SC8 and 2HRI and for AutoDock 4B18 were chosen as target and the binding energy, inhibition constant values of the 9 compounds are tabulated in Table 3. Consolidating the results of Docking, we selected best three compounds (Compound 11, 20 and 48) which were having better binding affinity and binding energy when compared with the standard PIPER compound. Hence these compounds can be used as a Scaffold to design more compounds of perylene diimides as well by using these scaffold, we can synthesize more compounds. These compounds may possess more potent and inhibitory activity for the telomerase enzyme which tends to apoptosis. Molecular modelling study helps to open future de novo modelling of new compound to treat cancer.

Table 3: Molecular Docking Study of Perylene ligands with the protein 4B18 using AutoDock 4.2 Software.

| S No | Compound | Conformation | Binding Energy | Ligand Efficiency | Inhibition Constant/ Units | No of Hydrogen Bonds | Hydrogen Contacts | Biological Activity |
|------|--------------|--------------|----------------|-------------------|----------------------------|----------------------|-------------------|---------------------|
| 1. | Compound _7 | 4th | -5.33 | -0.12 | 122.99 μ M | 1 | 7th:0:N38 | 5.87 |
| 2. | Compound _11 | 1st | -6.23 | -0.14 | 26.99 μ M | 1 | 4B18:A:GLN223:OE1 | 5.66 |
| 3. | Compound _20 | 2nd | -6.42 | -0.16 | 19.79 μ M | 1 | 11th:0:O18 | 5.59 |
| 4. | Compound _21 | 4th | -3.16 | -0.07 | 4.84 mM | 1 | 4B18:B:LYS236:N | 5.30 |
| 5. | Compound _30 | 2nd | -5.75 | -0.12 | 60.84 μ M | NIL | 20th: 0:O35 | 5.35 |
| 6. | Compound _33 | 4th | -1.62 | -0.04 | 64.72 mM | NIL | 4B18:A:TRP276:HE1 | 5.94 |
| 7. | Compound _42 | 4th | -2.89 | -0.06 | 7.63 mM | 1 | 21st: 0:O35 | 4.96 |
| 8. | Compound _48 | 10th | -6.59 | -0.04 | 67.8 mM | 1 | 4B18:B:SER229:OG | 4.84 |
| 9. | Compound _55 | 9th | -3.15 | 0.07 | 4.94 mM | 2 | NIL | 5.75 |
| 10. | PIPER | 4th | -5.59 | -0.12 | 80.55 μ M | 2 | NIL | 6.24 |

^aAll values are mean \pm SD (n=3)

ACKNOWLEDGEMENT

The authors are thankful to Vels University (VISTAS) and its management for providing research facilities and encouragement. The author is obliged to DBT - Government of India (BT/Bio-care/03/10047/2013-14) for providing financial assistance to carry out the research work.

CONFLICT OF INTEREST

The authors declare that they have no competing interests.

ABBREVIATIONS

PIPER: PIPER [N,N'-bis-(2-(1-piperidino)ethyl)-3,4,9,10-perylene tetracarboxylic acid diimide], **G4LDB:** Guanine Rich Ligand Database, **QSAR:** Quantitative Structure Activity Relationship, **VLife Science MDS-** V Life Sciences Molecular Design Suite, **PDB-** Protein Data Bank, **MGL-** Molecular Graphics Laboratory, **RCSB-** Research Collaboratory for Structural Bioinformatics.


REFERENCES

- Quantitative Structure Activity Relationship (QSAR) VLife Sciences Technologies Pvt Ltd. - VLife Product Documentation- Tutorial QSAR. - Obtaining 3D QSAR models by kNN - MFA method using VLife MDS.
- Quantitative Structure Activity Relationship (QSAR) VLife Sciences Technologies Pvt Ltd. - VLife Product Documentation- Tutorial QSAR.- Obtaining conventional QSAR model by Partial Least Squares Regression Method using VLife MDS.
- Hemalatha CN, Aanandhi MV. "G-Quadruplex Ligands as Stabilizer targeting Telomerase as Anti-Cancer Agents". Asian Journal of Pharmaceutical and Clinical Research. 2017;10:50-3.
- Hemalatha CN, Aanandhi MV. 3D QSAR and Docking study of Perylene-di imides analogues as potent apoptosis inducer and efficacious anticancer agent. Indian Drugs, Dec 2017 Issue, In Press. 2017.
- Blackburn EH. Telomeres and telomerase: Their mechanisms of action and the effects of altering their functions. FEBS Lett. 2005;579(4):859-62.
- Sanner MF. Python: A Programming Language for Software Integration and Development. J Mol Graphics Mod. 1999;17(1):57-61.
- Jain SV, Ghate M, Bhadoriya KS, Bari SB, Chaudhari A, Borse JS. 2D, 3D-QSAR and docking studies of 1, 2, 3-thiadiazole thioacetanilides analogues as potent HIV-1 non-nucleoside reverse transcriptase inhibitors. Organic and Medicinal Chemistry Letters. 2012;2(1):22. DOI: 10.1186/2191-2858-2-22.
- Franceschin M, Rossetti L, D'Ambrosio A, Schirripa S, Bianco A, Ortaggi G, *et al.* Natural and synthetic G-quadruplex interactive berberine derivatives. Bioorg Med Chem Lett. 2006;16(6):1707-11.
- Li Q, Xiang JF, Yang QF, Sun HX, Guan AJ, Tang YL. G4LDB: A database for discovering and studying G-quadruplex ligands. Nucleic Acids Research. 2012;41(D1):D1115-23.
- Salvati E. Telomerase damage induced by the G-Quadruplex ligand RHPS4 has an anti tumor effect. Journal of clinical investigation. 2007;117(11):3236-47.
- Rossetti L, Franceschin M, Schirripa S, Bianco A, Ortaggi G, Savino M. Selective interactions of perylene derivatives having different side chains with inter- and intramolecular G-Quadruplex DNA structures. A correlation with telomerase inhibition. Bioorganic and Medicinal Chemistry Letters. 2005;15(2):413-20.
- Laskowski RA, Mac Arthur MW, Moss DS, Thornton JM. PROCHECK - a program to check the stereochemical quality of protein structures. J App Cryst. 1993;26(2):283-91.
- Laskowski RA, Rullmann JA, Mac Arthur MW, Kaptein R, Thornton JM. AQUA and PROCHECK-NMR: Programs for checking the quality of protein structures solved by NMR. J Biomol NMR. 1996;8(4):477-86. [PubMed id: 9008363]

SUMMARY

- From QSAR 59 compounds are selected for Docking study. Docking has been done by G-Quadruplex Ligand Database (G4LDB). This is an online database which was having in built tools and performed by Open Babel 2.3.0 to predict the binding affinity with the targets. The targets (1LIH, 1NZM, 3CE5, 3SC8 and 2HRI) for the docking are selected based upon the literature survey and the selected compounds are docked. The results are visualized by Discovery studio Visualizer 4.1 Visualizer. From the results 9 compounds are selected and the results of these compounds are visualized using Discovery Studio 4.1 Visualizer. To narrow down the results, 2 best compounds are selected and the compounds are Compound 20 and Compound 48. The hydrogen bond interactions and binding free energy levels, pKi values are compared with the standard PIPER compound.
- From the results of G4LDB Database, the 9 compounds are docked with the specific protein (PDB ID: 4B18), a telomerase protein. Docking has been done by Auto Dock 4.2. Finally from the AutoDock results, Compound 11 shows good binding energy when compared with the standard PIPER compound. The study states as from the G4LDB Database 2 compounds possess good binding affinity, and from the Auto Dock Compound 11 possess good binding energy.
- The best potent compounds are studied to know the interactions with the nearby amino acids and the ligand interactions are visualized by online PROCHECK server.
- From the insilico studies, Compound 11, 20 and 48 has been showing best binding energy with respect to the target telomerase enzyme.

PICTORIAL ABSTRACT



About Authors



Ms. C.N.Hemalatha, M.Pharm., (Ph.D) is a Research Scholar carrying out Pharmaceutical Chemistry research at School of Pharmaceutical Sciences, Vels Institute of Science, Technology and Advanced Studies, VISTAS, Chennai.



Dr. M. Vijey Aanandhi: Is the Professor and Head of Department of Pharmaceutical Chemistry and Analysis at School of Pharmaceutical Sciences, Vels Institute of Science, Technology and Advanced Studies, VISTAS, Chennai.

Cite this article: Hemalatha CN and Muthkumar VA. Application of 3D QSAR and Docking Studies in Optimization of Perylene diimides as Anti Cancer Agent. Indian J of Pharmaceutical Education and Research. 2018;52(4):666-75.